

Altosid[®] Liquid Larvicide MOSQUITO GROWTH REGULATOR

PREVENTS EMERGENCE OF ADULT FLOODWATER MOSQUITOES

SPECIMEN LABEL

ACTIVE INGREDIENT:

(S)-Methoprene* 5%

OTHER INGREDIENTS: 95%

Total 100%

* CAS # 65733-16-6

Formulation contains 0.43 lb/gal (51.3 g/liter) active ingredient.

EPA Reg No. 2724-392

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE ADDITIONAL PRECAUTIONARY STATEMENTS

Because of the unique mode of action of **A.L.L.**[™], successful use requires familiarity with special techniques recommended for application timing and treatment evaluation. **See Guide to Product Application** or consult local Mosquito Abatement Agency.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS

CAUTION

Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling.

FIRST AID

If in eyes • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-248-7763 for emergency medical treatment information.

ENVIRONMENTAL HAZARDS

Do not contaminate water when disposing of equipment washwaters or rinsate.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

CHEMIGATION

Refer to supplemental labeling entitled "Guide to Product Application" for use directions for chemigation. Do not apply this product through any irrigation system unless the supplemental labeling on chemigation is followed.

MIXING AND HANDLING INSTRUCTIONS

1. SHAKE WELL BEFORE USING. **A.L.L.** may separate on standing and must be thoroughly agitated prior to dilution.
2. Do not mix with oil; use clean equipment.
3. Partially fill spray tank with water; then add the recommended amount of **A.L.L.**, agitate and complete filling. Mild agitation during application is desirable.
4. Spray solution should be used within 48 hours; always agitate before spraying.

APPLICATION INSTRUCTIONS

INTRODUCTION

A.L.L. must be applied to 2nd, 3rd, or 4th larval instars of floodwater mosquitoes to prevent adult emergence. Treated larvae continue normal development to the pupal stage where they die. This insect growth regulator has no effect when applied to pupae or adult mosquitoes. **A.L.L.** has sufficient field life to be effective at appropriate rates when applied to larval stages under varying field conditions. For further information, see Guide to Product Application.

METHODS OF APPLICATION

AERIAL

Use the prescribed amount of **A.L.L.** listed below in sufficient water to give complete coverage. One-half to 5 gallons of spray solution per acre is usually satisfactory. Do not apply when weather conditions favor drift from areas treated.

GROUND

Determine the average spray volume used per acre by individual operators and/or specific equipment. Mix **A.L.L.** in the appropriate volume of water to give the rate per acre as indicated below.

APPLICATION RATE

Apply 3 to 4 fl oz of **A.L.L.** per acre (219 to 293 ml/hectare) in water as directed.

APPLICATION SITES

PASTURES

A.L.L. may be applied after each flooding without removal of grazing livestock.

RICE

A.L.L. must be applied to 2nd, 3rd, and/or 4th instar larvae of mosquitoes found in rice, usually within 4 days after flooding. **A.L.L.** treatment may be repeated with each flooding.

INTERMITTENTLY FLOODED NONCROP AREAS

A.L.L. may be applied as directed above when flooding may result in floodwater mosquito hatch.

Typical sites include freshwater swamps and marshes, salt marshes, woodland pools and meadows, dredging spoil sites, drainage areas, waste treatment and settling ponds, ditches and other natural and manmade depressions.

CROP AREAS

A.L.L. may be applied to irrigated croplands after flooding to control mosquito emergence. Examples of such sites are: vineyards, rice fields (including wild rice), date palm orchards, fruit and nut orchards, and berry fields and bogs. Irrigated pastures may be treated after each flooding without the removal of grazing livestock.

DENSE VEGETATION OR CANOPY AREAS

Apply an **A.L.L.** sand or BIODAC mixture using standard granular dispersal equipment. For detailed preparation instructions, refer to Guide to Product Application.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

STORAGE

Store in cool place. Store product away from other pesticides, food, and feed. In case of leakage or spill, soak up with sand or another absorbent material.

CONTAINER DISPOSAL

Triple rinse or equivalent. Then offer for recycling or reconditioning or puncture and dispose of in a sanitary landfill, or incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

PESTICIDE DISPOSAL

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risks of use and handling of this material when such use and handling are contrary to label instructions.

Always read the label before using the product.

For information, call 1-800-248-7763 or visit our Web site: www.altosid.com

Wellmark

Wellmark International
Schaumburg, Illinois U.S.A.

ZOECON
Professional
Products

Zoecon® A Wellmark International Brand
A.L.L.™, ALTOSID® Liquid Larvicide Concentrate, and
ZOECON®, are trademarks of Wellmark International.

May, 2003
Schaumburg, IL